



66 ROSSTOWN ROAD, CARNEGIE, VICTORIA, 3163
 PHONE: (03) 9563 5422 FAX: (03) 9563 6944
 www.carnegievets.com.au

DR. KETAN JIVANJEE (BSc, BVSc, MRCVS)


DR. KELLY WILKINSON (BVSc Hons.)

Passion, Pride and Perseverance in Service and Care

EASY GUIDE PREVENTATIVE HEALTH CARE IN CATS (Melbourne)

VACCINATIONS

Feline Panleukopaenia
 Calicivirus
 Herpes
 Chlamydia
 Leukaemia
 FIV



- Initial **three(or four) course kitten vaccination** followed by
- **Annual** vaccination

There are two regimes for vaccination dependent on whether your cat is an indoor or outdoor cat:

	8 weeks	12 weeks	15-16 weeks	18 weeks
Indoor	F3	F4	F4	
Outdoor	F3	F6	F6	FIV

There are numerous preparations available for the control and eradication of fleas and the elimination of gastro-intestinal worms. You will most likely encounter differing opinions and advice on which products to use for your companion. Outlined below is your clinic's recommendations with respect to parasite control. The guideline is simple, convenient and most importantly effective.

FLEA TREATMENT

- We recommend the use of **Frontline^R Advantage^R** or **Revolution^R** on a monthly basis. These products are all effective in interrupting the flea life cycle and providing environmental control also.
- Please follow two basic rules when applying these products



Rule 1: Apply the entire contents of the product directly on the skin, by parting the coat at the back of the neck

Rule 2: Do not allow them to come into contact with water on their coat for at least 48 hours after application.

GASTRO-INTESTINAL WORMING

This is very important as some gastro-intestinal worms can be passed onto humans.

- We recommend the use of **Profender^R** or **Milbemax^R**
- Profender^R is applied to the skin. It should be done at 8 weeks, 12 weeks and then 3 monthly after that.
- Milbemax^R is a small tablet. It should be given at 6 weeks, 8 weeks, 12 weeks and then on a 3 monthly basis.

